

Epidemiology of Schistosomiasis in Damboa, Gaboru and Baga (IDP) Camps in Maiduguri, Borno State.

Babagana umar 1.,pukuma M. S 2.

- 1. Department of Basic sciences, YobeSstate College ofAgriculture Gujba, PMB 1104, Damaturu, Yobe State, Nigeria.**
- 2. Department of Zoology Moddibbo Adama University of Technology PMB 2076, Yola Adamawa State, Nigeria**

Abstract

Schistosomiasis also known as Bilharziasis or snail fever is a common intravascular trematode infection most common in developing regions of Africa and Asia. Five major species of parasitic trematodes of the family Schistosomatidae including *Schistosoma haematobium*, *Schistosoma intercalatum*, *Schistosoma japonicum*, *Schistosoma mansoni*, and *Schistosoma mekongi*. Over 200 million people, almost all of them in developing countries, suffered from Schistosomiasis which is associated with economic losses, and frequently interferes with development projects. The disease is endemic in most African countries where up to one-third of school age children may be actively infected. Hence this study was aimed to assess the distribution and pattern of Schistosomiasis. 600 urine sample each were examined from 600 persons were a clean, pre-labelled screw-capped plastic container were distributed with instructions to collect urine. The samples were immediately transported to diagnostic laboratory for examination. Sedimentation method was use to examine the urine sample. The overall number of people infected with *Schistosoma haematobium* was 162(27.0%). The results also shows that age group 6-15 years has the highest infection rates of 92(43.6%) male infected with *Schistosoma haematobium* while 18(21.7%) of female infected with *Schistosoma haematobium* respectively. Followed by age group 16-25 with 34(20.8%) number of male infected with *Schistosoma haematobium*, while 10(13.2%) of female infected with *Schistosoma haematobium* respectively. It is recommended that the control of snail intermediate host and the infective stage (*cercariae*) would in no doubt reduce the rate of transmission, thereby reducing the prevalence of infections. It was then concluded that *Schistosomiasis* among the study area was highly prevalent. Therefore, routine treatment, diagnosis and surveillance of the disease should be done by community-based organization to reduce the menace.

Key words: Helminthes; Parasites; *Schistosomiasis*; *Schistosoma haematobium*; Snails;

* Corresponding author. Tel.: 08030658483
E-mail address: bgumar2005@gmail.com

Introduction

Schistosomiasis also known as Bilharziasis or snail fever is a common intravascular trematode infection most common in developing regions of Africa and Asia (Pugh, et al., 2008). Five major species of parasitic trematodes of the family Schistosomatidae including *Schistosoma haematobium*, *Schistosoma intercalatum*, *Schistosoma japonicum*, *Schistosoma mansoni*, and *Schistosoma mekongi*, infect humans. Schistosomiasis, in which the bladder is affected, is caused by infection with *Schistosoma haematobium*, while intestinal Schistosomiasis is caused by *Schistosoma mansoni* both of which occur mainly in Africa. Over 200 million people, almost all of them in developing countries, suffer from Schistosomiasis, which can cause urinary obstruction, organ damage or destruction and death. At the same time, Schistosomiasis is associated with economic losses, and frequently interferes with development projects, particularly water resource development projects such as dams, irrigation schemes, planned and unplanned forestry (Gryseels, et al., 2007). In 1993, the World Health Organization (WHO) noted that the prevalence and intensity of the disease have been increased in areas undergoing water resource development, especially irrigation (WHO, 2013). The disease is endemic in most African countries where up to one-third of school age children may be actively infected although not always aware of their status (Chidozie, et al., 2008). Schistosomiasis is a neglected disease and very few studies have described its epidemiology.

Thus studies are needed to understand the epidemiology of these infections in order to implement measures necessary for their control in this region. The distribution of Schistosomiasis varies considerably with regions. In developing countries, the true epidemiological picture is not clear because of inadequate research on this infection despite its relevance in planning Schistosomiasis control in any locality (Nmorsi, et al., 2011). The most common method of diagnosis of Schistosomiasis in epidemiological surveys carried out in Africa is the identification of eggs in the stool for *Schistosoma mansoni* or in the urine for *Schistosoma haematobium*. Drug treatment is still the principal method of control and the drug of choice is praziquantel, however the degree of recovery from the infection depends on the extent of the damage caused by the infection. Single dose praziquantel (40 mg/kg) is effective in reducing prevalence and in curtailing the disease (King, et al., 2008).

Description of the Study Area

Maiduguri, is the capital city of Borno State, Nigeria. It is one of the 27 Local Government Areas of the state located in the northern part of the country between latitudes 11° 50" north, 13° 09" east. It is part of the Sahelian region and has an area of 543km². With an estimated population of 1,907,600 (NPC, 2006).

It is located at the central parts of the state and its shared common boundary with Jere, Konduga, Kaga and Magumeri local government area of the state. The study area perhaps represents a rainbow coalition of many tribes with Kanuri being the dominant tribe, others are Babur bura, Chibork, Marghi, Fulfulde, Gwoza and Shuwa people then, Fulani herdsmen and Igbo traders forming the minority.

The relief of the area provides uplands that give rise to streams and the vegetation is that of savannah. It occupies a fertile land which receives much rainfall and they use the land for growing of millet, sorghum, Maize, rice, cotton, okra and vegetables. There are many streams, ponds and few earth dams for domestic, recreational and agricultural purposes mostly in the dry season. Most of the inhabitants go to the streams that overflow their banks during the rainy season though some may dry up in the dry season, ponds and dams, especially children and teenagers to wade, swim and for fishing purposes, of which large proportion of these children become infected and re-infected (Cheesebrough, 2008).

The climate of area is close to sub-tropical (hot semi-arid) with the monthly temperature ranging 35°C and 47°C and a relative humidity of 38.4% to 63.4%, with august having the highest relative humidity. The annual rainfall is about 73.8mm to 193.2mm, and is recorded higher in august. The rainy season starts from June to September and the dry season from October to May. The dry, cold and dusty harmattan usually starts from November to February. (Ikusemoran, M. and Jimme, A.M. 2014).

Research Design

The distribution of Schistosomiasis was determined by selecting 3 IDP camps (Damboa camp, Gamboru camp and Baga camp). In each of the camp, 200 person were randomly selected cutting across all ages, giving a total number of 600 person involved. With the cooperation of camp official and different organization rendering assistance, the aim of the study was explained to the people in detail. This was to seek and obtain their consents. Mode and sources of transmission,

effect and control measures of the disease was emphasized to them. At the beginning of the study consent was obtained from the camp official of the study area.

Collection of Specimen

Information such as their Sex, Age, Occupation, and Source of water supply was recorded in a hard covered notebook. A clean, pre-labelled screw-capped plastic container were distributed with instructions to collect the first and last drops of mid-day (10.00am – 2.00pm) urine to suit the diurnal rhythm corresponding to the peak output of Schistosomal eggs (WHO, 2013; Cheesebrough, 2008). 10% formal saline was used to preserve the specimen. The samples were then transported immediately to state specialist Hospital Maiduguri diagnostic laboratory for examination between 30 minutes to 2 hours.

Urine analysis

20 milliliters of urine sample was collected from each person and allowed to stand for 30-60 minutes to the bottom of the plastic containers by ordinary sedimentation method. The supernatant was gently decanted until almost 10 milliliters was left. The 10 milliliters was mixed and turned inside a clean centrifuge tube, and centrifuged at 500 – 1000 revolution per minutes for 5 minutes. The supernatant was gently decanted off to leave only the deposits. Using a clean Pastuer pipette, a drop of the sediment was placed on a clean grease-free microscope slide and a cover slip was gently lowered on it, avoiding air bubbles. Then, it was viewed under x10 and x40 microscope objectives respectively, for the characteristics terminal-spine of *Schistosoma haematobium* eggs (Cheesebrough, 2008).

Chi-square test and percentage was used to analyze the result of the various parameters, such as the age groups, sex, and prevalence. Each parameter was calculated to determine whether or not an association exists between the parameter and the infection. Where $p < 0.05$ was considered significant. Using statistix8.0 version

Results

Table 1: shows the overall distribution of Schistosomiasis among the camps in relation to sex indicated that gamborou camp has the highest rate of 57 (28.5%) person infected with *Schistosoma haematobium*. Out of that 46 (33.1%) male infected while 11 (18.0%) female infected with *Schistosoma haematobium* respectively. Followed by бага camp has a total of 54 (27.0%) person infected with *Schistosoma haematobium*, out of that 44 (31.4%) male infected while 10 (16.7%) female infected with *Schistosoma haematobium* respectively. And lastly damboa camp has the

lowest rate of 51 (25.5%) person infected with *Schistosoma haematobium*. out of that 41 (29.1%) male infected with *Schistosoma haematobium* while 10 (16.9%) and 6 (10.2%) female infected with *Schistosoma haematobium* respectively.

Table 2: shows a prevalence of infection in relation to sex which indicate that out of 600 person examined, 420 and 180 were male and female respectively. A total of 162 (27.0%) person infected with *Schistosoma haematobium*. 131 (31.2%) male infected while 31 (17.2%) female with *Schistosoma haematobium* respectively. The results also shows that there was a significant difference in infection between the sexes where male has the highest rate of infection than the female.

Table 3: shows the distribution of Schistosomiasis among the study subjects in relation to age group, where it shows highest prevalence rate of infection among subjects belonging to age group 6-15 with 294 persons examined. Out of which 110 (37.4%) person infected with *Schistosoma haematobium*. Age group 36-45 has 22 person examined where 2 (9.1%) person infected with *Schistosoma haematobium*. Other age group were 16-25 with 239 person examined where 44 (18.4%) person infected with *Schistosoma haematobium*. And age group 26-35 has 44 number of person examined with 6 (13.6%) infected with *Schistosoma haematobium*. 46-above has no infection. Statistical analysis shows that there were significant difference in infection between the age group.

Table 1: Prevalence of Schistosomiasis among study subjects in the study camps in relation to sex.

Sex	Camps					
	Damboa No.examined	camp S. haematobium	Gamboru No.examined	camp S. haematobium	Baga No.examined	camp S. haematobium
Male	141	41(29.1%)	139	46 (33.1%)	140	44 (31.4%)
Female	59	10 (16.9%)	61	11 (18.0%)	60	10 (16.7%)
Total	200	51 (25.5%)	200	57 (28.5%)	200	54 (27.0%)

Table 2: Prevalence of Schistosomiasis in relation to sex.

	Number examined	Number (%) infected with S. haematobium
Male	420	131(31.2%)
Female	180	31 (17.2%)

Total	600	162 (27.0%)
Chi –Square (χ^2)		12.47**
Degrees of freedom		0.0004
P-value		1

Table 3: Distribution of Schistosomiasis among the study subjects in relation to age groups.

Age group	Number examined	Number (%) infected with S. haematobium
6-15	294	110 (37.4%)
16-25	239	44 (18.4%)
26-35	44	6 (13.6%)
36-45	22	2 (9.1%)
46-above	1	0 (0.0%)
Total	600	162 (27.0%)
Chi-square (χ^2)		33.06**
Degrees of freedom		0.0000
P-value		4

Table 4: Shows the prevalence of Schistosomiasis in the study area in relation to age group and sex. This table shows that age group 6-15 has the highest rate of 92 (43.6%) male infected while 18 (21.7%) female infected with Schistosoma haematobium respectively. Lower rate of 2 (13.3%) male infected while 10 (0.0%) female infected with Schistosoma haematobium was recorded in age group 36-45. Age group 16-25 has rate of 34 (20.0%) male infected while 10 (13.2%) female infected with Schistosoma haematobium respectively. 26-35 age group has rate of 3 (10.0%) male infected while 3 (21.4%) female infected with Schistosoma haematobium. And lastly 46-above had no male or female infected with Schistosoma haematobium respectively.

Table 5: Shows prevalence of Schistosomiasis among the three (3) camps in the study area. This table shows that in each of the camps two hundred (200) person were examined given a total of 600 person examined. Gamborou camp has highest prevalence rate of 57 (28.5%) person infected with Schistosoma haematobium. Followed by baga camp with prevalence rate of 54 (27.0%) persons infected with Schistosoma haematobium. And lastly damboa camp has a rate of 51 (25.5%) person infected with Schistosoma haematobium. Chi-square analysis shows that there is little or no significant difference in infection between the camps in relation to Schistosoma haematobium

Table 6: Shows the overall prevalence of *Schistosoma haematobium* and *Schistosoma mansoni* in relation to occupation among the study subjects in the three (3) IDP camps was 162(27.0%). The results also shows that pupils and students has the highest rate of 67(36.6%) and 69(22.9%) number infected with *Schistosoma haematobium* respectively. Others are housewife 11(31.4%), out of school children 9(17.3%), unemployed 5(21.7%) number infected with *Schistosoma haematobium* respectively. And only 1(100.0%) of farmer infected with *Schistosoma haematobium*. Chi-square analysis shows that there was significant difference between the infection of *Schistosoma haematobium*, and the people's occupation.

Table 4: Prevalence of Schistosomiasis in the study area in relation to age group and sex.

Age group	No. examined	Male Number(%) infected with <i>S. haematobium</i>	Female Number(%) infected with <i>S. haematobium</i>
6-15	294	92(43.6%)	18(21.7%)
16-25	239	34(20.8%)	10(13.2%)
26-35	44	3(10.0%)	3(21.4%)
36-45	22	2(13.3%)	0(0.0%)
46-above	1	0(0.0%)	0(0.0%)
Total	600	131(31.2%)	31(17.2%)

Table 5: Prevalence of Schistosomiasis among the camps in the study area.

Camps	No. examined	Number (%) infected with <i>S. haematobium</i>
Damboa camp	200	51 (25.5%)
Gamboru camp	200	57 (28.5%)
Baga camp	200	54 (27.0%)
Total	600	162 27.0%)
Chi-square (χ^2)		0.46*
Degrees of freedom		0.7959
P-value		2

Table 6: Prevalence of Schistosomiasis among the study subjects in relation to occupations.

Occupation	Number examined	Number (%) infected with <i>S. haematobium</i>
Pupils	183	67(36.6%)
Students	301	69(22.9%)
Housewife	35	11(31.4%)
Out of school children	52	9(17.3%)
Civil servant	5	0(0.0%)
Farmer	1	1(100.0%)
Unemployed	23	5(21.7%)
Total	600	162(27.0%)
Chi-square (χ^2)		16.67**
Degrees of freedom		0.0106
P-value		6

Discussion

The results of this study showed that Schistosomiasis was prevalent in the study area with overall prevalence of 162(27%). This results agrees with Nale et al., (2009) who reported a prevalence of 22.5% and this could be attributed to many outdoor activities engaged in infected water. The prevalence of infection in the camps could be attributed to closeness of the people from water bodies infected with snail intermediate host (Okon et al., 2007). Those live close to the water bodies or irrigation canals were more exposed and therefore more vulnerable to Schistosomiasis than those who lived further from the water Ugbomoiko et al., (2010), Abdullahi et al., (2011). Gaboru camp had 57(28.5%) of Schistosomiasis in urine because of their closeness to water bodies, followed by Baga camp 54(27.0%) and lastly the Damboa camp with 51(25.5%) Schistosomiasis respectively. This give a total of 162(27.0%) infected by *Schistosoma haematobium*.

It is clear from the results that the study area is endemic of Schistosomiasis. This results agreed with (Kiran and Muddasiru, 2014; Pukuma and Musa, 2007) Who reported a prevalence rate of urinary Schistosomiasis at (60.80%), and that of intestinal Schistosomiasis at (2.93%), which may

be attributed to water contact activities in the area. In relation to sex, the high infection rate observed in males than in females was also observed in other endemic areas as found by other authors (Ekejindu et al., 2002; Pukuma and Musa, 2007). This high prevalence in males than in females may be connected with the socio-cultural setup of the people of the study area. These people are predominantly Muslims, Hausa and Kanuri by tribe. Majority of the females are restricted to their houses therefore they have less contact with infested water compared to their male counterparts. Swimming and bathing in the open water bodies is also very uncommon among females in community. This is in line with the observation made by other authors (Bello et al., 2003).

The infection rates in the study area varied according to their ages, where age group 6-15 years had the highest infection rates of 92(43.6%) of male infected with *Schistosoma haematobium* while 18(21.7%) of female infected with *Schistosoma haematobium* respectively. Followed by age group 16-25 with 34(20.8%) number of male infected with *Schistosoma haematobium*, while 10(13.2%) female infected with *Schistosoma haematobium* respectively. This might be attributed to frequent water contact since those age groups engaged in activities that involved frequent contact with water. Nnoruka (2000) reported in Mayo Belwa, Adamawa state. A prevalence rate of *Schistosoma haematobium* in children between the ranges of 11-13 years. In Zuru Kebbi state, the highest prevalence was among age group 11-15 years. Joseph et al., (2010) and Akinboye et al., (2011) in their separate works showed higher prevalence of 15.0% in Maiduguri and 12.5% in Ibadan, respectively among school children of age group 12-15 years. However, results of this study agrees with Okoli et al., (2006) who reported the highest prevalence of 22.2% in the age group 21-30 years cohort in Ohaji/Egbema LGAs, Imo State Nigeria. The results of this study do not agree with Ombugadu, (2001) who reported peak prevalence of 40.2% and 28.6% in male and female between age group 21-25.

This results also showed that no infection rates was recorded in age group 46 above, this also agrees with Dawet et al., (2012) who reported no infection among age group 40- above, the low prevalence in aged people may be due to progressive increase in the level of naturally acquired immunity against Schistosomiasis and less contact with water.

Based on occupation, the overall study showed that pupils and students has the highest peak of 67(36.6%) and 69(22.9%) infected with *Schistosoma haematobium* respectively. followed by

housewives 11(31.4%), out of school children 9(17.3%), and unemployed 5(21.7%), *Schistosoma haematobium* respectively. And only 1(100.0%) of farmer infected with *Schistosoma haematobium*. This study does not agree with Pukuma et al., (2006) who reported a high prevalence of 38.9% among civil servant in Shelleng Town, Adamawa state. This could be associated with the fact that they can go to farms, ponds, streams for their domestic and recreational activities.

Conclusion and Recommendations

In conclusion, this study has indicated high prevalence of Schistosomiasis among the study area. Therefore, routine treatment, diagnosis and surveillance of the disease should be done by community-based organization to reduce the menace. The infection recorded could probably be due to reasons such as unhealthy environment, poverty, socio-cultural practices, lack of adequate health care facilities and ignorance. It has been observed that people in the study area were ignorant of the mode of transmission of this disease and proper sanitation which can increase contamination of the environment. The overall number of people infected with *Schistosoma haematobium* in the study area is 162(27.0%).

Prevalence of Schistosomiasis includes basically the destruction of intermediate snail host and infective stage (cercariae) with molluscicides and larvicides respectively. The control of snail intermediate host would in no doubt reduce the rate of transmission, thereby reduction in prevalence of infections. Workshops, seminars and control campaign programmed should be organized to train village heads who in turn educate their subjects on the mode of transmission, control strategies and dangers of the disease. Health education is a very effective means of improving knowledge about Schistosomiasis and the potential to reduce the prevalence of the diseases Jamda et al., (2007). Subjects in the study area should be educated by the local government health workers on proper means of waste disposal and construction of sanitary latrines in their homes so as to reduce the act of urinating or defecating in the open surroundings. The communities in the study area should embark on a monthly environmental sanitation. There should be provision of recreational centres in the communities to reduce the rate of contact with infected water, and village heads as well as opinion leaders should discourage on some of the socio-cultural practices (like urinating and bathing in stagnant or slow moving water) that may expose them to infection.

REFERENCE

- Abdullahi, M.K., Bassey, S.E. and Oyeyi, T.I. (2011). The Epidemiology of *Schistosoma haematobium* Infection in the 44 Local Government Area of Kano State, Nigeria. *Nigerian Journal of Parasitology*, **32**: 19 – 24.
- Akinboye, D.O., J.U. Ajisebetu, O. FAwole, O.M. Agbolade and O.O. (2011). Urinary Schistosomiasis: Water Contact Frequency and Infertility among Secondary School Students in Ibadan, Nigeria, *Nigerian Journal of Parasitology*, **32**: 129 – 134.
- Cheesebrough, M. (2008). *District Laboratory Practice in Tropical Countries*, 2nd Edition; P. 106 – 108.
- Chidozie, J., Patrick, G., Cletus, D.C., Reuben, O. (2007). Urinary Schistosomiasis among School Children in Ebonyi State, Nigeria, *International Journal of Laboratory Medicine*.
- Dawet, A., C.B. Benjamin, D.P. Yakubu (2012). Prevalence and Intensity of *Schistosoma haematobium* among Resident of Gwong and Kabong in Jos North Local Government Area, Plateau State, Nigeria. *International Journal of Tropical Medicine*, **7**(2) 69 – 73.
- Ekejindu IM, Ekejindu GOC, Andy A (2002). *Schistosoma haematobium* infection and nutritional status of residents in Azi-anam, a riverine area of Anmbara State, South-Eastern Nigeria. *Niger. J. Parasitol.* **23**:133-138.
- Gryseel, B., Polman, K., Clerinx, J., Kestens, L. (2007). Human Schistosomiasis. *Lancet*. **368**(9541): 1106 – 18.
- Ikusemoran, M. and jimme, A.M. (2014) A Decades Assessments of Maiduguri urban expansion (2002-2012): Geospatial Approach. *Global journal of geography, geo management* volume 14.
- Jamda, A.M., Ogbonna, C., Zoakah, I.A. and Daboer, J.C. (2007): Impact of health education on knowledge and practices of urinary Schistosomiasis among children in Martin village *journal of tropical medicine* **9**:21-27.
- Joseph, M.B., B. Gaji, T. Mohammed, M.M. Baba and I.B. Thilza (2010). Incidence of Schistosomiasis in Primary School Pupil with Particular Reference to *Schistosoma haematobium* in Maiduguri, *Researcher*, **2**: 31 – 36.
- King CH, Dangerfield-Cha M (2008). The unacknowledged impact of chronic Schistosomiasis. *Chronic Illn.* **4**: 65-79.
- Kiran S, Muddasiru D (2014). Epidemiology of schistosomiasis in school aged children in some riverine areas of Sokoto, Nigeria. *Niger. J. Parasitol.* **6**(6):197-201.
- Nale, V., Galadima, M. and Yakubu, S.E. (2009). Index of Potential Contamination of Urinary Schistosomiasis in Five Settlement near River Kubannai in Zaria, Nigeria. *Nigerian Journal of Parasitology* **24**: 95 – 101.

- Nmorsi OPG, Egwunyenga OA, Ukwandu NCD, Nwokolo NQ (2011) Urinary Schistosomiasis in a rural community in Edo state, Nigeria: Eosinophiluria as a diagnostic marker. *Afr. J. Biotechnol.* **4**: 183-186.
- Nnoruka, V.C. (2009). *Epidemiological Studies of Urinary Schistosomiasis*. Pub: Nigerian Journal of Parasitology, Vol. **21**, p. 21 – 23.
- Ombugadu, R.J. (2001) A Study of *Schistosoma haematobium* among the inhabitants of Udege community in Nassarawa State, Nigeria. *Biol., Environ. Sci., journal of tropical Medicine*, **8**:8184.
- Okoli, C.G., J.C. Anosike, and M.O.E. Iwuala (2006). Prevalence and Distribution of Urinary Schistosomiasis in Ohaji/Egbema Local Government Area of Imo State, Nigeria. *Journal Am. Science*, **2**:46 – 49.
- Okoli, C.G. (2007). Urinary Schistosomiasis in Cross River State, Nigeria. *Nigerian Journal of Parasitology* **17**:40 – 47.
- Okon, O.E.E., Udoutun, M.F., Nta, A.I., Etim, S.E., Abram, J.T. and Akpan, P.A. (2007). Prevalence of Urinary Schistosomiasis in Abini Community, Biase Local Government Area of Cross River State, Nigeria. *Nigerian Journal of Parasitology*. Vol. **28**(1): 28 – 31.
- Pugh RN, Gille HN (2008). Malumfashi endemic research project III Urinary Schistosomiasis; a longitudinal study. *Ann. Trop. Med Parasitol.* **72**: 271-482.
- Pukuma MS, Musa SP (2007). Prevalence of urinary schistosomiasis among residents of Waduku in Lamurde Local Government Area of Adamawa State Nigeria. *Niger. J. Parasitol.* **28**(2):65-68.
- Pukuma, S.M., M. Sale, H.L. Njila and M. Dibal (2006). A High Prevalence of 65 and 38.9% in Fishermen and Civil Servant Respectively was reported among School Pupils in Shelleng Town, Adamawa State, Nigeria. *Journal of Infectious Pest Disease Vector Management*, **7**:45 – 448.
- Ugbomoiko U.S., Ofoezie, I.E., Okoye, I.C. and Heukelback, J. (2010) Factors associated with urinary Schistosomiasis in two peri-urban communities in south-western Nigeria. *Annal tropical medical parasitology*, **104**:409-419.
- World health organization (WHO), 2013. Schistosomiasis, fact sheet no 115. Available at www.who.int/mediacentre/factsheet/fs115/ sheet.