

Comparing The Effectiveness of Normal Saline, Chlorhexidine-Setrimid and Propylbetaine Polihexanide as Wound Washing Solution On Pseudomonas Aeruginosa Contaminated Wound

Muhammad Irzal Wahab^{a*}, Sitti Rizaliyana^b, Lobredia Zarasade^b

^aichalwahab@gmail.com

Department of Plastic Reconstructive and Aesthetic Surgery Faculty of Medicine Airlangga University, Surabaya, Indonesia

Abstract

Background: The wound healing process is closely related to wound care techniques which consist of cleansing, debridement and dressings. In infected wounds, washing with a wound washing solution is expected to reduce the colony of germs and to eradicate the infection. One of the important pathogens in wound is Pseudomonas aeruginosa. In this study we will compare the use of normal saline, chlorhexidine-cetrimide, and propylbetaine-polyhexanide as wound washing solutions for wounds contaminated with Pseudomonas aeruginosa.

Methods: Twenty-seven mice were divided into 3 groups: normal saline, chlorhexidine-cetrimide, and propylbetaine-polyhexanide group. The wound was made on the mice's back then the Pseudomonas aeruginosa germ was inoculated on the wound. Four hours later, the wound was irrigated with the wound washing solution according to its group. Tissue culture will be taken 18 hours after washing.

Result : Propylbetaine-Polihexanide is the most effective wound washing solution on Pseudomonas aeruginosa contaminated wound compare to Normal Saline and Chlorhexidine-Cetrimide. Although, there is no significant difference in the number of Pseudomonas aeruginosa bacteria on the wound bed after washing with Normal Saline, Chlorhexidine-Cetrimide, and Propylbetaine-Polihexanide.

Keyword : Wound Solution; normal saline; chlorhexidine-setrimid; propylbetaine-polyhexanide; Pseudomonas aeruginosa; white mice

1. Introduction

Wound healing process is closely related to wound management or techniques which consist of cleansing, debridement and dressings. The wound care components above support each other in the wound healing process, however, cleansing is a measure of the success of wound care and is a standard wound care practice that is always practiced by professional health practitioners both in Indonesia and abroad (Baranoski & Ayello, 2012).

In the United States, Pseudomonas aeruginosa is the most common pathogen in hospitals and the second most common bacterial pathogen in patients with ventilator acquired pneumonia (Gellatly & Hancock, 2013). Research conducted by Asadullah (2015) on patients with nosocomial pneumonia at Dr. Soetomo Hospital Surabaya for the period January 1, 2011 - March 30, 2012, showed that the most pathogenic bacteria causing nosocomial pneumonia were Pseudomonas aeruginosa (Asadullah, Isbandiyah, & Sri Ardila, 2012).

In Dr. Soetomo Hospital, the standard antiseptic for wound management is chlorhexidine-cetrimide, more commonly known as Savlon®. While in several other hospitals using wound washing solution like normal saline (NaCl 0,9%) and propylbetaine - polyhexanide (Prontosan®)

2. Methods

Twenty-seven mice were divided into 3 groups: normal saline, chlorhexidine-cetrimide, and prophyllbetaine-polyhexanide group. A full-thickness wound size of 1x1 cm was made on the back's mouse using a knife and scissors. *Pseudomonas aeruginosa* for contaminated wound in mice was obtained from positive culture isolates results of Dr. Soetomo Hospital's patients. Bacteria were incubated and cultured on standard media, then placed on the wound and covered with transparent dressing.

Four hours after application of *Pseudomonas aeruginosa*, each wound was irrigated using NaCl 0.9%, Savlon® 1:30, and Prontosan® according to the treatment group, using a 20cc syringe within 30 seconds, to the entire surface of the wound. Wound cleansing was implemented with irrigation alone, without any deep wiping of the wound, to prove the efficacy of the washing fluid. Tissue culture were performed on all parts of the wound 18 hours after washing the wound (Picture 1).

In calculating the number of colonies, score 1 is given if there is no germ growth, score 2 if the growth of germs is less than 10^5 colonies, and score 3 is given if it grows more than 10^5 colonies. Calculation of the number of colonies using colony forming units per cm^2 (CFU / cm^2).




Fig .1. Research Framework

3. Result

Table 1. Score of colonies number calculation

Treatment	Score					
	1		2		3	
Washing with Normal Saline (0,9% NaCl)	0	0%	0	0%	9	100%
Washing with Chlorhexidine - Cetrimide 1:30	0	0%	2	22,2%	7	77,8%
Washing with Propylbetaine-Polihexanide	2	22,2%	2	22,2%	5	55,6%

On the table 1, washing treatment group with normal saline shows point 3 has a percentage of 100%. The washing treatment group with chlorhexidine - cetrimide 1:30 showed that point 2 had a percentage of 22.2% and point 3 had a percentage of 77.8%. From the propylbetaine-polihexanide washing treatment group showed that point 1 showed a presentation of 22.2%, point 2 showed a presentation of 22.2%, and point 3 showed a presentation of 55.6%.

The results obtained were then analyzed statistically using a nonparametric test, namely the Kruskal Wallis.

Table 2. The Kruskal Wallis

Kruskal Wallis H	df	Asymp. Sig.
5,272	2	0,072

On table 2, it informs the difference in the number of *Pseudomonas aeruginosa* bacteria in wound beds after washing with normal saline, chlorhexidine-cetrimide, and propylbetaine-polihexanide resulted in an H test statistic of 5.272 with a significance value of 0.072. It can be seen that the significance value > significant alpha (5% or 0.05), means there is no significant difference in the number of *Pseudomonas aeruginosa* bacteria on the wound bed after washing with Normal Saline, Chlorhexidine-Cetrimide, and Propylbetaine-Polihexanide.

4. Discussion

4.1 Colonization

Wounds which were made in the operating room and prepared by asepsis procedures were sterile wounds. Meanwhile wound that were not obtained that way such as by accident, the wound is called a contaminated wound. Contaminated wounds if left and have passed the contamination time will become an infected wound. Infected wounds are called dirty wounds. Contamination wounds, no matter how contaminated with external dirt, can still be cleaned during treatment and are not called dirty wounds, they are still contaminated wounds (Djohansjah, 2018).

Contamination time is the time when the bacteria in the wound are still in the wound room, not yet infecting the tissue so that it is easy to clean. Generally, it lasts around 6-8 hours. For areas of the face and neck that are rich in vascularization, this time can be longer than other parts, around 12-18 hours. (Djohansjah, 2018)

4.2 *Pseudomonas aeruginosa*

Pseudomonas aeruginosa is a nosocomial pathogen most often responsible for infections in patients with cystic fibrosis, bronchiectasis, neutropenia, AIDS, and / or burns and patients with metabolic, hematological, or malignant diseases. This bacterium is mainly responsible for ventilator-related pneumonia, infections after surgery, and urinary tract infections, and sepsis in patients in intensive care units. Most strains are generally resistant to many antibiotics including imipenem and meropenem; therefore, pathogens are usually problematic in therapy (Yetkin, 2006).

The ability of *Pseudomonas aeruginosa* to form biofilms makes *Pseudomonas aeruginosa* a feared pathogen because of the combined effects of various intrinsic virulence factors, and its resistance to antimicrobial drugs, and due to *Pseudomonas* contamination can cause death (Hamijaya, 2014).

Physical characteristics of wounds infected with *Pseudomonas aeruginosa* are the presence of a lot of suppuration, turquoise in color and gives color to the bandage, pungent and disturbing smell; tend to infect other patients in the same ward; difficult to heal and very damaging to the wound; and can be followed by diarrhea due to septicemia, as well as weakness and anemia in patients (Turner & Richardson, 1917).

4.3 Normal Saline (NaCl 0,9%)

Normal saline is a widely recommended wound washing fluid, because it is known to be compatible with human tissue. This fluid does not cause damage to new tissue and does not affect the function of fibroblasts and keratinocytes in wound healing. Its effectiveness in preventing infection has not been established (Moscato, 1998; Penn-Barwell, 2012).

Many cleaning solutions have shown safe and effective results, while others can damage and destroy cells that are important for the healing process. Normal saline is considered the most appropriate and preferred cleaning fluid because this solution is non-toxic, isotonic. fluid that does not damage healing tissue (Davis, 2017).

4.4 Chlorhexidine - Cetrimide 1:30 (Savlon®)

One of the popular antiseptic ingredients is from the biguanide group, namely chlorhexidine. Chlorhexidine has extensive antimicrobial activity and is less irritating to the skin, but it kills organisms slower than alcohol, and decreases its activity if there is organic material and alkaline atmosphere in the wound. Chlorhexidine is the active ingredient most widely used not only in various antiseptic and disinfectant products but also in handwashing, preservative and mouthwash preparations. Cetrimide as a surfactant is added to chlorhexidine and has a non-specific destructive effect on bacterial membranes (McDonnell & Russell, 1999).

4.5 Propylbetaine-polihexanide (Prontosan®)

Prontosan® wound solution, is a wound cleanser made of pure water and two main ingredients: polyhexamethylene, antimicrobial agent (0.1%) betaine, surfactant (0.1%) (Bradbury, 2011).

Although water can be used as a wound cleanser, and has not been seen to increase the risk of infection or delay healing, the use of specially designed wound cleaning agents may have the potential to improve clinical outcomes through their additional wound cleaning modalities. Evidence appears that the combination of polyhexamethylene with surfactants (betaine) has an increased ability to penetrate hard-to-remove layers, removing bacteria and biofilms from wounds (Bradbury, 2011).

The results of this study confirm previous studies that can prove the effectiveness of Propylbetaine-Polihexanide against *Pseudomonas aeruginosa*. A previous study by Minnich (2012) showed a reduction in the in vitro bacterial load in 12 of the 13 micro-organisms commonly found in wounds. As with all in vitro studies, the observed results may not translate into clinical benefit or represent the full range of situations seen in clinical practice. However, this study provides objective data on the antimicrobial activity of the product in vitro, adding that the affected patients gave the impression of clinical effectiveness of 0.1% polyhexanide / 0.1% betaine for wound cleansing (Davis, 2017).

In the case of irrigation for wound washing, the decrease in the rate of bacterial proliferation due to irrigation of wounds in the wound with Propylbetaine-Polihexanide solution, gave rise to evidence that the combination of polyhexamethylene with a surfactant (betaine) has an increased ability to penetrate difficult-to-remove layers, removing bacteria and biofilms from the surface. wounds (Bradbury, 2011).

5. Conclusion

Propylbetaine-Polihexanide is the most effective wound washing solution on *Pseudomonas aeruginosa* contaminated wound compare to Normal Saline and Chlorhexidine-Cetrimide, although there is no significant difference in statistic.

Reference

- Asadullah, Isbandiyah, & Sri Ardila, 2012. Pola bakteri penyebab pneumonia nosokomial di RS Dr Soetomo Surabaya periode Januari 2011 - Maret 2012. *Saintika Medika Journal*, 11(1): pp. 64 – 67. <https://doi.org/10.22219/sm.v11i1.4197>
- Atiyeh, B.S., Dibo, S.A., & Hayek, S.N., 2009. Wound cleansing, topical antiseptics and wound healing. *International Wound Journal*, 6(6): pp. 420 – 430. <https://doi.org/10.1111/j.1742-481X.2009.00639.x>. PMID: 20051094.
- Baranoski, S., & Ayello, E.A., 2012. *Wound Care Essentials: Practice Principles*, 3rd ed. Philadelphia: Lippincott Williams & Wilkins.
- Bellingeri, A., Falciani, F., Traspedini, P., Moscatell, A., Russo, A., Tino, G., Chiari, P., & Pedhetti, A., 2016. Effect of a wound cleansing solution on wound bed preparation and inflammation in chronic wounds: a single-blind RCT. *Journal of Wound Care*, 25(3): pp. 160 – 168. <https://doi.org/10.12968/jowc.2016.25.3.160>. PMID: 26947697.
- Bradbury, S., & Fletcher, J., 2011. Prontosan made easy. *Wounds International*;2(2): pp. 1–6.
- Davidson, J.M., & DiPietro, L., 2008. The wound-healing process. In: A. Veves, J.M. Giurini, & F.W. Langerfo, eds. *The Diabetic Foot: Medical and Surgical Management*, 2nd ed., Totowa: Humana Press, pp. 59 – 82.
- Davis, S.C., Harding, A., Gil, J., Parajon, F., Valdes, J., Solis, M., & Higa, A., 2017. Effectiveness of a polyhexanide irrigation solution on methicillin-resistant *Staphylococcus aureus* biofilms in a porcine wound model. *International Wound Journal*, 14(6): pp. 937 – 944. <https://doi.org/10.1111/iwj.12734>. PMID: 28266133.
- Delignette-Muller, M.L., 1998. Relation between the generation time and the lag time of bacterial growth kinetics. *International Journal of Food Microbiology*, 43(1-2): pp. 97–104. [https://doi.org/10.1016/s0168-1605\(98\)00100-7](https://doi.org/10.1016/s0168-1605(98)00100-7). PMID: 9761343.
- Fernandez, R.S., Griffiths, R., & Ussia, C., 2007. Water for wound cleansing. *International Journal of Evidence-Based Healthcare*, 5(3): pp. 305 – 323. <https://doi.org/10.1111/j.1479-6988.2007.00068.x>.

- Gellatly, S.L., & Hancock, R.E.W., 2013. *Pseudomonas aeruginosa*: new insights into pathogenesis and host defenses. *Pathogens and Disease*, pp: 159 – 173. <https://doi.org/10.1111/2049-632X.12033>. PMID: 23620179.
- Grey, J.E., Enoch, S., & Harding, K.G., 2006. Wound assessment. In: J.E. Grey & K.G. Harding, eds. *ABC of Wound Healing*. Cardiff: Wiley-Blackwell, pp. 1 – 4.
- Gurtner, G.C., 2007. Wound healing, normal and abnormal. In: C.H.M. Thorne, R.W. Beasley, S.J. Aston, S.P. Bartlett, & G.C. Gurtner, eds., *Grabb and Smith's Plastic Surgery*, 6th ed., Philadelphia: Lippincott Williams & Wilkins, pp. 15 – 22
- Hamijaya, L., Prihatiningsih, & Widiastuti, M.G., 2014. Perbedaan daya anti bakteri tetrachlorodecaoxide, povidon iodine, dan hidrogen peroksida (H₂O₂) terhadap bakteri *pseudomonas aeruginosa* secara invitro. *Jurnal Kedokteran Gigi*, 5(4), pp. 329 – 335.
- James, W.D., Berger, T.G., Elston, D.M., & Odom, R.B., 2006. Skin: basic structure and function. In: W.D. James, T.G. Berger, & D.M. Elston, *Andrews' Diseases of the Skin: Clinical Dermatology*, 10th ed., Philadelphia: Elsevier, pp. 1 – 32.
- Kanno, E., Kawakami, K., Ritsu, M., Ishii, K., Tanno, H., Toriyabe, S., Imai, Y., Maruyama, R., & Tachi, M., 2011. Wound healing in skin promoted by inoculation with *Pseudomonas aeruginosa* PAO₁: The critical role of tumor necrosis factor- α secreted from infiltrating neutrophils. *Wound Repair and Regeneration*, 19(5): pp. 608–621. <https://doi.org/10.1111/j.1524-475X.2011.00721.x>. PMID: 22092799.
- Kanno, E., Tanno, H., Suzuki, A., Kamimatsuno, R., & Tachi, M., 2016. Reconsideration of iodine in wound irrigation: the effects on *Pseudomonas aeruginosa* biofilm formation. *Journal of Wound Care*, 25(6): pp. 335–339. <https://doi.org/10.12968/jowc.2016.25.6.335>. PMID: 27286666.
- Kanno, E., Toriyabe, S., Zhang, L., Imai, Y., & Tachi, M., 2010. Biofilm formation on rat skin wounds by *Pseudomonas aeruginosa* carrying the green fluorescent protein gene. *Experimental Dermatology*, 19(2): pp. 154–156. <https://doi.org/10.1111/j.1600-0625.2009.00931.x>. PMID: 19650869.
- Kolarsick, P.A.J., Kolarsick, M.A., & Goodwin, C., 2011. Anatomy and physiology of the skin. *Journal of the Dermatology Nurses' Association*, 3(4): pp. 203 – 213. <https://doi.org/10.1097/JDN.0b013e3182274a98>.
- Köljal, S., Naaber, P., & Mikelsaar, M., 2002. Antibiotic resistance as an indicator of bacterial chlorhexidine susceptibility. *Journal of Hospital Infection*, 51(2): pp. 106 – 113. <https://doi.org/10.1053/jhin.2002.1204>. PMID: 12090797.
- Kryger, Z.B., & Sisco, M., 2007. *Practical Plastic Surgery*, 1st ed., Texas: Landes Bioscience.
- Lima, C.C., Pereir, A.P., Silva, J.R., Oliveira, L.S., Resck, M.C.C., Grechi, C.O., Bernardes, M.T.C.P., Olimpio, F., Sanos, A.M.M., Incarpi, E.K., & Garcia, J.A.D., 2009. Ascorbic acid for the healing of skin wounds in rats. *Brazilian Journal of Biology*, 69(4): pp. 1195 – 1201. <https://doi.org/10.1590/S1519-69842009000500026>.
- Maier, R.M., 2009. Bacterial growth. In: R.M. Maier, L.L. Pepper, & C.P. Gerba, eds., *Environmental Biology*, 2nd ed., San Diego: Academic Press, pp. 37–54.
- Marzoeki, D., 1993. *Luka dan Perawatannya*, 1st ed., Surabaya: Universitas Airlangga Press, pp. 9 – 36.
- McDonnell, G., & Russell, A.D., 1999. Antiseptics and disinfectants: activity, action, and resistance. *Clinical Microbiology Reviews*, 12(1): pp. 147 – 179. PMID: 9880479. PMCID: PMC88911.
- Moscato, R., Mayrose, J., Fincher, L., & Jehle D, 1998. Comparison of normal saline with tap water for wound irrigation. *American Journal of Emergency Medicine*, 16(4): pp. 379 – 381. [https://doi.org/10.1016/S0735-6757\(98\)90133-4](https://doi.org/10.1016/S0735-6757(98)90133-4).

- Murray, P.R., Rosenthal, K.S., & Pfaller, M.A., 2016. Medical Microbiology, 8th ed., Philadelphia: Elsevier.
- Nagoba, B.S., Selkar, S.P., Wadher, B.J., & Gandhi, R.C., 2013. Acetic acid treatment of pseudomonal wound infections - a review. *Journal of Infection and Public Health*, 6(6): pp. 410 – 415. <https://doi.org/10.1016/j.jiph.2013.05.005>. PMID: 23999348.
- Nurman, M., 2017. Perbandingan efektifitas madu + NaCl 0,9% dengan NaCl 0,9 % saja terhadap penyembuhan luka gangren pada pasien Diabetes Mellitus tipe II di wilayah kerja puskesmas Bangkinang Kota tahun 2015. *Jurnal Ners*, 7(1): pp. 53 – 62.
- Penn-Barwell, J.G., Murray, C.K., & Wenke, J.C., 2012. Comparison of the antimicrobial effect of chlorhexidine and saline for irrigating a contaminated open fracture model. *Journal of Orthopaedic Trauma*, 26(12): pp. 728 – 732. <https://doi.org/10.1097/BOT.0b013e31826c19c4>. PMID: 22874117.
- Perdanakusuma, D., 2008. Anatomi Fisiologi Kulit dan Penyembuhan Luka. Surabaya: Airlangga University Press.
- Queirós, P., Santos, E., Apostolo, J., Cardoso, D., Cunha, M., & Rodrigues, M., 2014. The effectiveness of cleansing solutions for wound treatment: a systematic review. *JB I Database of Systematic Reviews and Implementation Reports*, 12(10), pp. 121 – 151.
- Regan, M.C., Kirk, S.J., Wasserkrug, H.L., & Barbul, A., 1991. The wound environment as a regulator of fibroblast phenotype. *Journal of Surgical Research*, 50(5), pp. 442 – 448.
- Rossolini, G.M., & Mantengoli, E., 2005. Treatment and control of severe infections caused by multiresistant *Pseudomonas aeruginosa*. *Clinical Microbiology and Infection*, 11(Supplement 4), pp. 17 – 32. <https://doi.org/10.1111/j.1469-0691.2005.01161.x>. PMID: 15953020.
- Salton, M.R.J., 1968. Lytic agents, cell permeability, and monolayer penetrability. *The Journal of General Physiology*, 52(1): pp. 227s – 252s. PMID: 4913700.
- Sarabashi, S., 2010, Anatomy of the skin. In: S. Sarabashi, ed., *Principles and Practice of Burn Care*, 1st ed, New Delhi: Jaypee Brothers Medical Publisher, pp. 18 – 24.
- Taylor, K., 1916. Treatment of *Bacillus pyocyaneus* infection. *JAMA*, LXVII(22), pp. 1598 – 1599. <https://doi.org/10.1001/jama.1916.02590220040013>.
- Thomas, G.W., Rael, L.T., Bar-Or, R., Shimonkevitz, R., Mains, C.W., Sione, D.S., Craun, M.L., Bar-Or, D., 2009. Mechanisms of delayed wound healing by commonly used antiseptics. *The Journal of trauma*, 66(1): pp. 82 – 90. <https://doi.org/10.1097/TA.0b013e31818b146d>. PMID: 19131809.
- Turner, P., & Richardson, G., 1917. The treatment of wounds infected with *Bacillus pyocyaneus*. *British Medical Journal*, 2(2961): pp. 421 – 423. <https://doi.org/10.1136/bmj.2.2961.421>. PMID: 20768751. PMCID: PMC2349206.
- Werner, S., & Grose R, 2003. Regulation of wound healing by growth factors and cytokines. *Physiological reviews*, 83(3): pp. 835 – 870. <https://doi.org/10.1152/physrev.2003.83.3.835>. PMID: 12843410.
- Yetkin, G., Baris, O., Cicek, A., Kuzucu, C., & Durmaz, R., 2006. Clinical, microbiologic, and epidemiologic characteristics of *Pseudomonas aeruginosa* infections in a University Hospital, Malatya, Turkey. *American Journal of Infection Control*, 34(4): pp. 188 – 192. <https://doi.org/10.1016/j.ajic.2005.11.010>.
- Young, A., & McNaught, C.M., 2011. The physiology of wound healing. *Surgery*, 29(10): pp. 475-479. <https://doi.org/10.1016/j.mpsur.2011.06.011>